

Affiliative and disciplinary behavior of human handlers during play with their dog affects cortisol concentrations in opposite directions

Zsuzsanna Horváth*, Antal Dóka, Ádám Miklósi

Department of Ethology, Institute of Biology, Eötvös Loránd University Faculty of Science, Budapest, Hungary

Received 29 August 2007; revised 31 January 2008; accepted 1 February 2008

Available online 15 February 2008

Abstract

It has been shown that cortisol concentrations change characteristically in the course of agonistic interactions; our aim was to find out how a playful situation may affect concentrations of this hormone in the saliva. We studied dogs' behavior and the changes of cortisol concentrations in a play situation, where the dogs played with their handler for 3 min with a tug toy. In this experiment working dogs were divided into two groups by the type of their work, namely police dogs and border guard dogs. We found that the cortisol concentrations of old police dogs significantly increased, while the adult border guard dogs' hormone levels decreased, which shows that playing with the handler has an effect on both groups, but interestingly this effect was opposite. Behavior analysis showed differences only in the behavior of the human handlers during the play sessions, while the behavior analysis did not reveal significant differences in the two groups of dogs, except that old border guard dogs generally needed more time to begin playing than old police dogs. During the play sessions police officers were mainly disciplining their dogs, while the border guards were truly playing with them (including affiliative and affectionate behavior). Our results are in accordance with those of recent studies, which show that behaviors associated with control, authority or aggression increase cortisol concentrations, while play and affiliative behavior decrease cortisol levels.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Play; Stress; Cortisol; Communicative signals; Working dogs

Introduction

Play is a characteristic behavior of young animals (Bekoff, 2001), which is composed of action patterns that are used in various contexts, such as exploration, manipulation, locomotion, predation, fighting and mating (Hol et al., 1994), or combinations of these (Loizos, 1966). Although its evolutionary origin is not clear, many hypothesized functions of play activity have been proposed including its promotional effect on physical and social skills (Byers and Walker, 1995; Drea et al., 1996; Dugatkin and Bekoff, 2003; Holmes, 1995; Soderquist and Serena, 2000).

Play is often used as an indicator of well being (for a review see e.g. Boissy et al., 2007) as animals exposed to traumatic stimulation during development, or live in impoverished environments, show reduced play activity. If we define stress as the non-specific response of the body to any demand for change (Selye, 1950), namely as an increase in HPA axis activity caused by any physical, psychological or environmental stimulus (e.g. Sapolsky et al., 2000), then it seems that environmental stress can have an inhibitory effect on the expression of play behavior. However, there could also be an opposite relationship between play and stress. Some recent studies have shown that play activity may contribute to the reduction of stress (Arelis, 2006). Rats stimulated by novel objects increase their play activity and at the same time show reduced anxiety (Darwish et al., 2001), and in chimpanzees the intensity of playing increases before feeding times which are usually associated with high levels of social tension (Palagi et al., 2004). These latter authors assume that this behavior

* Corresponding author. Department of Ethology, Institute of Biology, Eötvös Loránd University Faculty of Science, Pázmány P. s. 1/C 1117 Budapest, Hungary. Fax: +36 1 381 2180.

E-mail address: horvath.zsuzsanna@gmail.com (Z. Horváth).

might have a 'preventive' effect to reduce subsequent social stress elicited by the feeding situation. Humans also utilize play routinely to familiarize children with novel situations including research settings.

It is often stated that in comparison to their wolf (*Canis lupus*) ancestors, the dog (*Canis familiaris*) is a very playful species, since dogs play levels remain high throughout their life (Bekoff, 1972; Lorenz, 1950). Dogs are valued members of human social groups partly as a result of their eagerness to engage in inter-specific play (Bekoff, 1972; Fagen, 1981; Mitchell and Thompson, 1993; Rooney et al., 2001; Russell, 1936), and many owners spend a considerable amount of their time playing with their dogs (Hart, 1995).

As playful interaction between dogs and humans represents a natural activity for both species (see above), in the present study we used this behavior to investigate the relationship between play and stress by measuring saliva cortisol concentrations before and after dog–human playing interaction.

In social situations other than play, it has been shown that human behavior toward the dog has significant effects on its physiological state (e.g. change of cortisol concentrations, heart rate variability). For example, the presence of humans in a shelter (Beerda et al., 1997; Hennessy et al., 1998; Tuber et al., 1996) may be an effective means of reducing the cortisol response of dogs. In a recent study, Jones and Josephs (2006) found that after agility competitions there is a significant correlation between humans' punitive behaviors (physically pushing the dog and yelling at it) and the increase in dogs' cortisol concentrations in the losing teams. We also recently reported that threatening behavior shown by humans resulted in increasing cortisol concentrations in dogs (Horváth et al., 2007).

Thus in this study we hypothesized that the actual motor and communicative aspects of behavior of the human partners during play affect playing experience and inner state of the dogs. Since pilot observations showed that there is a variation in the behavior of working dog handlers toward their dog (policemen and border guards, see below), this offered a good opportunity for testing our hypothesis. In our experiments we asked handlers to interact in a playful manner with their dog. To reveal how dogs respond to such a situation, we simultaneously gathered behavioral data from both dogs and handlers and measured the saliva cortisol concentrations of dogs before and after the interaction. The experimental protocol was designed to include situations such as playful struggle with the handler (tug-of-war game), but also cooperative behaviors like the retrieval of the toy or giving the toy up to the handler. Using this method we aimed to document the changes of cortisol concentrations that take place during the playful episodes in dogs, in parallel to the actual behavior of the handlers.

Materials and methods

Subjects

The police dogs were purchased by the Hungarian National Police Training School for Police Dog Handlers (Dunakeszi, Hungary). Dogs were acquired between 1 and 3 years of age. Dogs were tested physically (i.e. for hip dysplasia) and behaviorally (i.e. reaction to gun shot, bite work). Individuals were

purchased only if they did not show signs of hip dysplasia and fear of gun shot. Thereafter they participated in a 12 week training course together with their handlers. During this course the dogs were trained for guarding and obedience.

Dogs that participated in the present study were purchased between 1997 and 2003, and performed patrol service with their handlers on the streets for a minimum of 1 year. All 84 subjects were male German Shepherd Dogs. The dogs' age ranged from 2 to 11 years (mean age±SD: 6.91±2.19 years) and the subjects were categorized following Studzinski et al. (2006): *Adult dogs* were 2–7 years old (43 individuals; mean age±SD: 5.18±1.45 years); *Old dogs* were 8–11 years old (41 individuals; mean age±SD: 8.76±0.94 years). We also divided the dogs into two work groups ('occupation'): *Police dogs* (53 individuals; mean age±SD: 7.26±2.05 years; 25 adult dogs (mean age±SD: 5.52±1.44 years) and 28 old dogs (mean age±SD: 8.82±0.94 years)), and *Border Guard dogs* (31 individuals; mean age±SD: 6.42±2.32 years; 18 adult dogs (mean age±SD: 4.77±1.47 years) and 13 old dogs (mean age±SD: 8.61±0.96 years)). Dogs of both groups were selected according to the same criteria initially and they also took part in identical training programs. The only differences were that some dogs were handled by policemen and the others by border guards and also in the nature of their work after the training period. Police dogs and their handlers are assigned to patrol duty on the streets, while border guard dogs and their handlers work the borders of Hungary in the countryside. Policemen and border guards were also selected with similar criteria. Eighty two of the handlers were men and 2 were women.

All procedures were approved by the Ethical Committee of Eötvös Loránd University, Department of Ethology and conducted in accordance with the Hungarian State Health and Medical Service (ÁNTSZ). There is a standing agreement with the Hungarian Police Force that permits testing their working dogs.

Date and premises

The experiments were carried out in 2005 and 2006 at the Hungarian National Police Training School for Police Dog Handlers (Dunakeszi, Hungary). The police dogs that were tested, participated in a special 2 week training course with their handlers during the time of testing. The experiments were conducted in an empty room (10 m long×10 m wide×5 m high), where only the experimenter (who was also the camera person), the handler and the dog were present during the test. The experimental room was familiar for the subjects (some of the training exercises take place at this location), as it has been shown that introduction into a novel environment enhances HPA activity in the dog (e.g. Beerda et al., 1997). All salivary samples were taken between 9 a.m. and 3 p.m. (e.g. Dreschel and Granger, 2005; Jones and Josephs, 2006). The dogs were in their kennels resting for 30 min before starting the test.

Procedure

Dogs played with their handler for 3 min in a similar manner to that which is described by Rooney and Bradshaw (2002). The handler could use a rag or a tug toy (a piece of thick rope with two knots on both ends, 20 cm long) for inducing play behavior in the dog. The handlers were instructed to play as intensively as possible with the dog, and to adjust their behavior to the dog's reactions. Before the test, the handler was asked to leave the dog's toys or food items outside the experimental room. The handlers were asked to encourage the dog during the entire trial, even if the dog displayed only slight or no inclination to play. We imposed few restrictions in the type of play; however the handlers were instructed to execute the following acts at least once during a play session: (1) they had to throw the object and encourage the dog to bring it back; (2) they had to try to take the object from the dog's mouth.

Measurement of saliva cortisol concentrations

Saliva samples were collected from the dogs before the play session and 20 min after the end of the play session (see also Beerda et al., 1998; Dreschel and Granger, 2005; Jones and Josephs, 2006; Vincent and Michell, 1992). Substances to stimulate saliva flow were not used. The saliva was collected with cotton swabs by the handlers near to the location of the experiment. While the dog was standing still, the handler placed the swab into the mouth of the dog and held it there until it absorbed the greatest amount of saliva possible (lasting from 30 to 60 s). The soaked cotton swabs were temporarily

stored on dry ice in numbered Eppendorf tubes. For long term storage the saliva samples were kept in a deep freezer (-80°C). Before analysis the tubes were warmed up to room temperature. The saliva was removed from the cotton swabs by centrifugation (3000 rpm for 15 min) using special centrifuge tubes with filters (Corning Spin-X; Sigma-Aldrich Kft., Budapest, Hungary). After separation the saliva samples were analyzed for cortisol concentrations using a highly sensitive (from 0.003 to 3.0 $\mu\text{g}/\text{dl}$) enzyme immunoassay kit from Salimetrics (State College, PA, USA); the intra- and inter-assay coef-

ficients of variation as provided by the manufacturer, are below 10% and 15% respectively (Salimetrics, 2005). The procedures were performed on the basis of the protocol provided by Salimetrics. Before calculating concentrations, log transformations were used to establish normal distributions. All analyses used the log-transformed hormone values. However, non-transformed data are reported in the figures to facilitate interpretation. All participating dogs provided the required amount of saliva and all cortisol measurements could be used for the statistical analysis.

Table 1

Behavioral variables coded in dogs and handlers, where 0 points are given (score 0) in case of the behavior that is least favorable from the point of view of our cooperative situation, while 2 points (score 2) are given in case of the most favorable behavior

	Behavioral variables	Score 0	Score 1	Score 2
DOGS	'Motivation'	There is no physical contact between the dog and the play object (e.g. the dog never holds or chews the object during the session)	The dog makes physical contact (with his mouth or paw) with the object at least once	There are two or more physical contacts made between the dog and the object
	'Playfulness'	The dog is passive, pays no attention to play object	The dog shows interest (looks at, contacts, etc.) in playing with the object at least one time	The dog pays attention to the play object during the entire session or more than one time
	'Willingness to retrieve'	The dog never brings the object to the handler or there is no physical contact between the dog and the object (e.g. the dog never holds or chews the toy during the session)	The dog makes steps to move towards the handler with the object in its mouth, but the handler cannot get the object without approaching the dog	The dog usually brings the object back to the handler
	'Possessivity' (this behavior variable was not coded when the dog was given Score 0 at 'Motivation' and/or 'Playfulness')	The handler is unable to take the object from the dog during the session even with force	There are visible signs of resistance when the handler tries to take the object or the dog shows avoidance with the object in its mouth, but finally the handler can take it from the dog	The handler can take the object from the dog's mouth (without any visible signs of force)
	'Aggression' (signs of aggression after Kim et al. (2006), Kroll et al. (2004), McLeod (1996), Netto and Planta (1997) and Pal et al. (1998): tail is not tucked, the ears are forward or ambivalent, stiff body posture; growling, barking, snarling, showing the teeth)	No signs of aggression can be observed during the play session	The dog shows at least one of the behaviors described during the play session	
	'Fear' (signs of fear after Dreschel and Granger (2005) and Kroll et al. (2004): the dog's tail is tucked between its hind legs, hides behind the human's leg, backs away or retreats, ears are pinned or drawn back or down)	No signs of fear can be observed during the play session	The dog shows at least one of the behaviors described during the play session	
	'Latency of starting to play' (sec; from the handler's first call to play till the dog takes the tug toy)			
HANDLERS	'Control commands' (number of verbal orders issued by the handler during the 3 min session, e.g. using words of command: sit — ül, stand — áll, lay — fekszik, heel — lábhoz, listen — figyelj)	Does not occur	Occurs between 1 and 5 times	Occurs 6 or more times
	'Sound signal' (number of non-verbal sound signals used by the handler during the 3 min session to attract the dogs' attention, e.g. whistle, clapping, etc.)	Does not occur	Occurs between 1 and 5 times	Occurs 6 or more times
	'Enthusiasm of handler'	The handler does not laugh or smile during the play session	The handler is cheerful during the play session	
	'Praising' (handler praises the dog or speaks to the dog kindly, in a high pitched or fluctuating voice)	Does not occur	Occurs between 1 and 5 times	Occurs 6 or more times
	'Petting head and body' (handler extends hand to touch or pet the dog anywhere on the head or body)	Does not occur	Occurs between 1 and 5 times	Occurs 6 or more times

Analysis of behavior

The behavior of the dogs was videotaped and analyzed later. From the video recordings, 11 different behaviors were scored based on a subjective assessment of the intensity of the behaviors (frequency), with one variable measured as an absolute latency. The behavioral variables are in Table 1.

The subsequent statistical analysis was based on the behavior scoring recorded by an experienced person (Zs. H.), who also coded the entire sample. However 20% of the recordings were also coded by another naïve, independent observer. For data coded by both observers, Kappa coefficients were calculated to measure inter-observer agreement and relatively high values were calculated in all cases. The values of Kappa coefficients are as follows: ‘motivation’: 1.0; ‘playfulness’: 1.0; ‘willingness to retrieve’: 1.0; ‘possessivity’: 0.91; ‘latency of starting to play’: 0.75; ‘control commands’: 0.83; ‘sound signals’: 0.85; ‘enthusiasm of handler’: 0.77; ‘praising’: 1.0; ‘petting head and body’: 0.85.

Questionnaire information

After the test, the handlers were asked to fill in a questionnaire which included questions about the dog and the dog–human relationship. Based on this questionnaire we could gain information about the handlers (age, experience with this dog in years), the living arrangements of the dogs, how much time the dog–handler pair spends patrolling and how often the handlers play with the dogs. The exact questions regarding this information are presented in Table 2.

Statistical analysis

Shapiro–Wilk’s test was used to assess whether or not the sample is consistent with a specified distribution function.

Cortisol concentrations were log transformed for analyses. The salivary cortisol data were analyzed using ANOVAs for repeated measures followed by Bonferroni’s post-hoc test. Three-way ANOVAs with repeated measures were carried out with ‘occupation’ (two levels: police dogs, border guard dogs) and ‘age’ (two levels: adult, old) as between-subjects factors, and ‘sampling’ (two levels: baseline, 20 min post) as within-subjects factor. We used paired *t*-tests for comparing the ‘baseline’ and ‘20 min post episode’ cortisol concentrations

Table 2
Summary table for the questionnaires filled in by the handlers

Answers to the questionnaire	Police	Border guard
Where is the dog kept (dog’s accommodation (%))?		
Block of flats	0	0
Flat with common garden	5.66	3.22
Family house with garden	62.26	96.77
Farm	3.77	0
Kennel at the police station	28.3	0
How many times per week does the handler play with the dog? (%)		
Never	3.77	3.22
Once/week	11.32	12.9
Three times/week	16.98	38.7
Five times/week	41.5	35.48
Every day	26.41	9.67
How much time is spent with the dog in patrol duties? (mean±SD hours/day)	5.87±2.87	9.26±3.83*

Accommodation and playing with the dog are exclusive categories (tick boxes on the questionnaire) and duration of the duty had to be judged by the handler. The bold numbers signify that more border guards kept their dogs at the family house with a garden than policemen, while more policemen kept their dogs in the kennel at the police station than border guards. Also border guards played three times per week with their dogs which was more than policemen, while more policemen played with their dogs daily which was more than border guards. The star (*) shows that border guards spend significantly more time on duty with their dogs than policemen do.




Fig. 1. The cortisol concentrations after the test of adult border guard dogs was significantly lower than before, while the cortisol concentration increased significantly in old police dogs (the connecting lines indicate the interactions). Post hoc analysis showed that the baseline cortisol concentrations were significantly higher in adult border guard dogs than in police dogs. Non-transformed data are presented as median±quartiles in µg/dl. Sample size of the subgroups: APD: *N*=25 (mean age±SD: 5.52±1.44 years); OPD: *N*=28 (mean age±SD: 8.82±0.94 years); ABGD: *N*=18 (mean age±SD: 4.77±1.47 years); OBGD: *N*=13 (mean age±SD: 8.61±0.96 years). BGD — border guard dogs; PD — police dogs; A — adult dogs; O — old dogs.

within work groups, and independent sample *t*-test for comparing the ‘baseline’ between working groups.

The behavior of police and border guard dogs, adult and old dogs, policemen and border guards, was calculated using Mann–Whitney U-tests. Spearman’s Rank correlation was used to search for relationships between non-parametric behavior variables measured in the case of policemen and border guards. We used also Spearman’s Rank correlation to search for relationships between humans’ behavioral variables and dogs’ cortisol concentrations. We used the Chi-Square test for analyzing the questionnaire data provided by the policemen and border guards.

Statistical analyses were performed using SPSS (Version 13.0). A significance level of 0.05 was adopted throughout.

Results

The analysis of salivary cortisol concentrations

The three-way ANOVA revealed that in general, the ‘occupation’ did not have a significant effect on the cortisol concentrations ($F(1,80)=0.18$; $p=0.67$), nor was there an effect of ‘sampling’ ($F(1,80)=0.03$; $p=0.8$), or ‘age’ of the dog ($F(1,80)=0.15$; $p=0.7$). Conversely, we found significant interaction between ‘sampling’ and the ‘occupation’ ($F(1,80)=9.0$; $p=0.004$). The cortisol concentrations after the test were significantly higher than before the test in police dogs ($t=-2.93$; $p=0.005$), while cortisol concentrations after the test were significantly lower than before the test in border guard dogs ($t=2.04$; $p=0.05$). When comparing the ‘baseline’ cortisol levels of police and border guard dogs, we found no significant differences between the two groups ($t=1.94$; $df=82$; $p=0.055$). We also found no significant interaction between ‘sampling’

and ‘age’ groups ($F(1,80)=0.14$; $p=0.7$). The interaction was significant between work (‘occupation’) and ‘age’ groups ($F(1,80)=4.47$; $p=0.038$), indicating that adult border guard dogs had higher basal cortisol concentrations than the adult police dogs, while the old police dogs had higher post-playing cortisol concentrations than the old border guard dogs (Fig. 1). The three-way interaction between ‘sampling’, the work groups (‘occupation’) and the ‘age’ groups, was not significant ($F(1,80)=0.28$; $p=0.6$).

The analysis of behavior

Behavior of dogs

When comparing the behavior of all police and border guard dogs, we found no significant differences in any of the behavioral variables (Table 3).

Comparisons of the behavior of adult police and border guard dogs did not reveal any significant differences in any of the behavioral variables. However, old border guard dogs generally needed more time to begin playing, than old police dogs (‘latency of play’: $Z=-2.2$; $p=0.038$).

In the case of police dogs, we found no significant differences in behavior between adult and old dogs. However, old border guard dogs generally needed more time to begin playing than younger border guard dogs ($Z=-2.46$; $p=0.022$) (Fig. 2).

Handlers' behavior

The police officers used more ‘control commands’ in the play situation ($Z=-2.8$; $p=0.005$), while border guards ‘pet’




Fig. 2. We found significant (*) differences in the latency of starting to play between adult and old dogs within the groups of border guard dogs (□ white boxes), and between old border guard dogs (□ white box) and old police dogs (■ grey filled box), as well. The old border guard dogs needed more time to begin playing. Sample size of the subgroups: APD: $N=25$ (mean age±SD: $5.52±1.44$ years); OPD: $N=28$ (mean age±SD: $8.82±0.94$ years); ABGD: $N=18$ (mean age±SD: $4.77±1.47$ years); OBGD: $N=13$ (mean age±SD: $8.61±0.96$ years). Data are presented as median±quartiles and differences are considered statistically significant if $p<0.05$.

the dog more often ($Z=-3.7$; $p<0.001$), and generally showed more ‘enthusiasm’ in the play situation ($Z=-2.35$; $p=0.018$) (Fig. 3).

There was no difference between adult and old dogs in either of the behavioral variables measured, and both policemen (P) and border guards (BG) showed similar behavior toward their dog’s, independent from their dog’s age (Table 4).

We found positive correlation between ‘control commands’ and ‘praising’ ($r_s=0.349$; $p=0.009$) in policemen. The policemen gave a command and if the dogs obeyed it, they praised

Table 3
Statistical results of behavioral variables in the case of police and border guard dogs

Behavioral variables	Statistical results				
	PD vs. BGD	Adult PD vs. adult BGD	Old PD vs. old BGD	Adult PD vs. old PD	Adult BGD vs. old BGD
Motivation	$Z=-1.05$, $p=0.293$	$Z=-0.49$, $p=0.62$	$Z=-1.49$, $p=0.13$	$Z=-0.72$, $p=0.47$	$Z=-1.51$, $p=0.13$
Playfulness	$Z=-1.94$, $p=0.052$	$Z=-1.21$, $p=0.22$	$Z=-1.78$, $p=0.07$	$Z=-0.86$, $p=0.39$	$Z=-1.32$, $p=0.18$
Willingness to retrieve	$Z=-0.65$, $p=0.511$	$Z=-0.58$, $p=0.56$	$Z=-1.7$, $p=0.08$	$Z=-0.22$, $p=0.83$	$Z=-1.90$, $p=0.06$
Possessivity	$Z=-0.30$, $p=0.75$	$Z=-0.28$, $p=0.77$	$Z=-0.53$, $p=0.59$	$Z=-0.73$, $p=0.47$	$Z=-1.09$, $p=0.27$
Latency of starting to play	$Z=-1.9$, $p=0.057$	$Z=-0.88$, $p=0.38$	$Z=-2.19$, $p=0.03^*$	$Z=-1.76$, $p=0.08$	$Z=-2.46$, $p=0.02^{**}$
Aggression	–	–	–	–	–
Fear	–	–	–	–	–

The stars for the significance show that old border guard dogs needed significantly (*) more time to begin playing, than old police dogs; and old border guard dogs needed significantly (**) more time to begin playing than younger border guard dogs. We have not mentioned the results concerning fear and aggression, as fear was not coded in any of the subjects (–), while aggressive behavior was shown by only one subject (–). BGD — border guard dogs; PD — police dogs.




Fig. 3. The policemen (■ grey filled boxes) use significantly (*) more ‘control commands’ with their dogs (□ white boxes), while the border guards pet their dogs significantly (*) more. Data are presented as median±quartiles and differences are considered statistically significant if $p<0.05$.

Table 4
Statistical results of behavioral variables in the case of policemen (P) and border guards (BG)

Behavioral variables	Statistical results	
	P	BG
Control commands	$Z=-0.02; p=0.18$	$Z=-1.66; p=0.09$
Sound signals	$Z=-1.44; p=0.15$	$Z=-1.13; p=0.26$
Enthusiasm of handler	$Z=-0.32; p=0.75$	$Z=-1.2; p=0.23$
Praising	$Z=-0.13; p=0.89$	$Z=-0.89; p=0.37$
Petting head and body	$Z=-0.99; p=0.32$	$Z=-1.0; p=0.31$

their dogs similarly as they did during training. In the police dyads we found a weak negative correlation between ‘enthusiasm of handler’ and ‘latency of starting to play’ of the dog ($r_s=-0.302; p=0.025$), that is, if the handler was more cheerful, the playing of the dog began sooner. The border guards that were more enthusiastic (‘enthusiasm of handler’), tended to pet (‘petting head and body’) their dogs more often than the others ($r_s=0.413; p=0.019$).

Relationship between cortisol concentrations and behavior

In case of the police dogs, no significant correlations were found between ‘baseline’ cortisol concentrations and the dogs’ behavioral variables. However, we found a slight positive correlation between dogs’ ‘post playing’ cortisol concentrations and the amount of ‘control commands’ used by policemen ($r_s=0.272; p=0.049$).

In border guard dogs we found a positive correlation between ‘baseline’ cortisol concentrations and ‘motivation’ ($r_s=0.515; p=0.003$), and negative correlation between ‘baseline’ levels and ‘latency of starting to play’ ($r_s=-0.428; p=0.014$). Border guard dogs with high ‘baseline’ cortisol concentrations were more motivated to play than others and they started to play immediately. In addition, we found negative correlation between dogs’ ‘post playing’ cortisol concentrations and the ‘frequency of physical praising’ ($r_s=-0.37; p=0.041$).

Questionnaire information

There were no significant differences between the policemen and border guards in their age ($Z=-0.022; p=0.982$), or in their experience with their dog ($Z=-0.252; p=0.801$). We found significant differences in living arrangements, time spent in service and time spent weekly with play. The border guards only kept their dogs at home (‘dog’s accommodation’: $\text{Chi}^2=13.29; \text{df}=3; p=0.004$), while policemen often kept their dogs in a kennel at the police station. Border guards spent more time with their dogs in service (‘patrol duties’: $Z=-3.619; p<0.001$), in contrast to policemen. However, policemen reported that they play more with their dogs ($Z=-2.305; p=0.021$) than border guards, and we found a negative correlation between ‘time in patrol’ and ‘time spent with play’ ($r_s=-0.365; p=0.001$).

Discussion

The aim of this study was to examine the effect of human communicative and social behavior on the inner state (measured in terms of saliva cortisol concentrations) of dogs in a play situation. To this end, dogs and their handlers were observed in a playful interaction, before and after which we measured changes of the dogs’ cortisol concentrations.

We found that on the whole, short term play interaction with the handler did not significantly change the cortisol concentrations of the dogs at a population level. This was in contrast to our previous study in which we observed an overall increase in cortisol concentration after being exposed to an approach by a threatening human (Horváth et al., 2007). However, important insights emerged when we compared the results of our two groups, police and border guard dogs, in which the direction of change in the cortisol concentration was the opposite. Moreover, the age of the dogs seemed to also have an influence, as in adult border guard dogs, cortisol concentration decreased after play, whereas in the old police dogs, the cortisol concentration increased by the end of the play session.

We also found a relationship between border guard dogs’ baseline cortisol concentrations and motivation and latency of starting to play, namely border guard dogs with higher baseline levels are more motivated to play with the handler than those with a lower levels, as they generally start to play immediately. Notably, in contrast to the changes in cortisol concentrations, the play behavior of police dogs was similar to play of the border guard dogs (i.e. they appeared just as playful and motivated as border guard dogs). Although the behavioral analysis of the play behavior may have not revealed initial differences in the motivation, we believe there were different causal factors behind the play behavior of the dogs. That is, police dogs may have executed playful behavior as part of a training exercise (‘they were commanded to play’), while border guard dogs may have played spontaneously with their handler. As a result, the social play may have mediated an inner state which was associated with lower concentrations of cortisol.

The analysis revealed that the policemen continually disciplined (‘controlled’) their dogs and used sound signals to gain their dogs’ attention during play sessions, while border guards were more empathetic and enthusiastic, and also pet and praised their dogs more often. The positive correlation between enthusiasm of handler and dogs’ latency of starting to play in the case of policemen, also suggested that the more enthusiasm shown by handlers, the sooner their dogs started to play. Thus we suppose that the differences in behavior, mood and motivation of the handlers not only influenced their dogs’ motivation and behavior during the play session (O’Farrell, 1997), but also had an effect on the post-play cortisol concentrations, which is in accordance with the findings of previous studies (e.g. Coppola et al., 2006; Jones and Josephs, 2006; Tuber et al., 1996).

Accordingly, the disciplinary behavior of policemen resulted in higher cortisol concentrations in the case of old police dogs, while the friendlier attitude and petting by the border guards, reduced cortisol concentrations in the case of adult border guard

dogs. Police dogs might have assessed this situation as a training session due to the behavior of the handlers, rather than play time. Our results are similar to conclusions of Jones and Josephs (2006), who have revealed a relationship between larger increased cortisol concentrations and punitive behavior and lesser increased cortisol concentrations and affiliative behavior, respectively after an agility competition.

It is important to stress that other factors might have also contributed to this effect, such as the age of the dogs, previous keeping conditions and the established relationship between dogs and their handler.

In the case of age, we found that adult border guard dogs with higher baseline cortisol concentrations are more motivated to play than old ones, which in general is in agreement with age-related decrease in playing activity in dogs (e.g. Rooney et al., 2000). In our case, the more affiliative behavior of the handler contributed to the reduction of cortisol concentrations in the adult dogs, suggesting a calming effect of social interaction. In contrast, the more disciplinary attitude of the policemen affected the cortisol concentrations in old dogs in the opposite direction. A recent meta-analysis found greater cortisol response to a challenge in older people (Otte et al., 2005) and we have reported similar observation in old police dogs as a response to the approach of a threatening human (Horváth et al., 2007). Notably, the behavior of the handlers was similar toward adult and old dogs, therefore this cannot explain the present findings.

According to the handlers (questionnaire), many of the police dogs live in kennels, while all the border guards take their dogs home after work. In the course of the present training course, all of the dogs spent their day in kennels, which might have been a more stressful experience for the border guard dogs, as police dogs are used to living in such quarters. This difference could have contributed to the different reaction of the dogs to the play situation.

Furthermore, both previously mentioned factors could have also influenced the nature of the actual social relationship between dogs and handlers, which might in some form, mirror intraspecies relationships in dogs or wolves. In this case, the larger social distance in terms of dominance could also influence the calming effect of the interaction, partly because dominants may use more forceful and physically challenging behavioral actions during the interactions, whereas partners having a similar social status may induce interactions (including play) by using more communicative signals including signals for play invitation (e.g. play bow).

In summary, police dogs playing with their handlers showed some 'similarity to' experience obtained during general training. This could be partly attributed to the behavior of policemen, but other factors cannot be excluded. It follows that despite the session's playfulness, the interaction with police handlers increased cortisol concentrations in the dogs, suggesting increased level of stress. Contrasting processes have been revealed in border guard dogs where we found support for the stress reducing effect of play.

Further studies may reveal whether dogs living with families might be also affected by the play behavior of their owner. It seems that affiliative communicative signals during play might

indeed contribute to stress reduction (at least in terms of decreased cortisol concentrations) supporting earlier findings (Arelis, 2006; Palagi et al., 2004). Thus regular play with dogs could contribute to their well being in general and also reduction of stress in certain situations.

Acknowledgments

This study was supported by the Hungarian Academy of Sciences (F 226/98) EU FP-6 NEST012787 and an OTKA grant (T049615). The authors are grateful to the Hungarian National Police Training School for Police Dog Handlers (Dunakeszi, Hungary) for their cooperation and to the all participating policemen; special thanks to Pál Marsi police colonel and Ferenc Suszter police major. We would like to express our gratitude to Professor Imre Oláh, head of the Developmental Biology and Immunology Lab (Semmelweis University) for his cooperation. We would like to thank Professor Sergio M. Pellis, Canadian Centre for Behavioural Neuroscience for his valuable help concerning the literature. We thank Dorottya Ujfalussy for her valuable comments on previous versions of the manuscript. We would like to express our appreciation to Celeste R. West-Pongrácz for being kind enough to correct the language of our manuscript as a native English speaker. We are most grateful for the supportive comments of two anonymous referees.

References

- Arelis, C.L., 2006. Stress and the power of play. A Thesis Submitted to the School of Graduate Studies of the University of Lethbridge in Partial Fulfillment of the Requirements for the Degree. Department of Neuroscience, University of Lethbridge, Lethbridge, Alberta, Canada.
- Beerden, B., Schilder, M.B.H., van Hooff, J.A.R.A.M., de Vries, H.W., 1997. Manifestations of chronic and acute stress in dogs. *Appl. Anim. Behav. Sci.* 52, 307–319.
- Beerden, B., Schilder, M.B.H., van Hooff, J.A.R.A.M., de Vries, H.W., Mol, J.A., 1998. Chronic stress in dogs subjected to social and spatial restriction. II. Hormonal and immunological responses. *Physiol. Behav.* 66, 243–254.
- Bekoff, M., 1972. The development of social interaction, play, and meta-communication in mammals: an ethological perspective. *Q. Rev. Biol.* 47, 412–434.
- Bekoff, M., 2001. Social play behaviour. *J. Conscious. Stud.* 8, 81–90.
- Boissy, A., Manteuffel, G., Jensen, M.B., Moe, R.O., Spruijt, B., Keeling, L.J., Winckler, C., Forkman, B., Dimitrov, I., Langbein, J., Bakken, M., Veissier, I., Aubert, A., 2007. Assessment of positive emotions in animals to improve their welfare. *Physiol. Behav.* 92 (3), 375–397.
- Byers, J.A., Walker, C., 1995. Refining the motor training hypothesis for the evolution of play. *Am. Nat.* 146, 25–40.
- Coppola, C.L., Grandin, T., Enns, R.M., 2006. Human interaction and cortisol: can human contact reduce stress for shelter dog? *Physiol. Behav.* 87, 537–541.
- Darwish, M., Korányi, L., Nyakas, C., Almeida, O.F.X., 2001. Exposure to a novel stimulus reduces anxiety level in adult and aging rats. *Physiol. Behav.* 72, 403–407.
- Drea, C.M., Hawk, J.E., Glickman, S.E., 1996. Aggression decreases as play emerges in infant spotted hyenas: preparation for joining the clan. *Anim. Behav.* 51, 1323–1336.
- Dreschel, N.A., Granger, D.A., 2005. Physiological and behavioural reactivity to stress in thunderstorm-phobic dogs and their caregivers. *Appl. Anim. Behav. Sci.* 95, 153–168.
- Dugatkin, L.A., Bekoff, M., 2003. Play and the evolution of fairness: a game theory model. *Behav. Processes* 60, 209–214.
- Fagen, R., 1981. *Animal Play Behavior*. Oxford University Press, New York.

- Hart, L.A., 1995. Dogs as companions: a review of the relationship. In: Serpell, J. (Ed.), *The Domestic Dog: Its Evolution, Behaviour and Interactions with People*. Cambridge University Press, Cambridge, pp. 161–179.
- Hennessy, M.B., Williams, M.T., Miller, D.D., Douglas, C.W., Voith, V.L., 1998. Influence of male and female petters on plasma cortisol and behaviour: can human interaction reduce the stress of dogs in a public animal shelter? *Appl. Anim. Behav. Sci.* 61, 63–77.
- Hol, T., Koolhaas, J.M., Spruijt, B.M., 1994. Consequences of short-term isolation after weaning on later adult behavioural and neuroendocrine reaction to social stress. *Behav. Pharmacol.* 5, 88–89.
- Holmes, W.G., 1995. The ontogeny of littermate preferences in juvenile golden-mantled ground squirrels: effects of rearing and relatedness. *Anim. Behav.* 50, 309–322.
- Horváth, Z., Igyártó, B.-Z., Magyar, A., Miklósi, Á., 2007. Three different coping styles in police dogs exposed to a short-term challenge. *Horm. Behav.* 52, 621–630.
- Jones, A.C., Josephs, R.A., 2006. Hormonal interactions between man and the domestic dog. *Horm. Behav.* 50 (3), 393–400.
- Kroll, T.L., Houpt, K.A., Erb, H.N., 2004. The use of novel stimuli as indicators of aggressive behavior in dogs. *J. Am. Anim. Hosp. Assoc.* 40 (13), 19.
- Loizos, C., 1966. Play in mammals. *Symp. Zool. Soc. Lond.* 18, 1–9.
- Lorenz, K., 1950. *Man meets dog*. Routledge Classics, Great Britain.
- Mitchell, R.W., Thompson, N.S., 1993. Familiarity and the rarity of deception: Two theories and their relevance to play between dogs (*Canis familiaris*) and humans (*Homo sapiens*). *J. Comp. Psychol.* 107 (3), 291–300.
- O'Farrell, V., 1997. Owner attitudes and dog behaviour problems. *Appl. Anim. Behav. Sci.* 52, 205–213.
- Otte, C., Hart, S., Neylan, T.C., Marmar, C.R., Yaffe, K., Mohr, D.C., 2005. A meta-analysis of cortisol response to challenge in human aging: importance of gender. *Psychoneuroendocrinology* 30, 80–91.
- Pal, S.K., Ghosh, B., Roy, S., 1998. Agonistic behaviour of free-ranging dogs (*Canis familiaris*) in relation to season, sex and age. *Appl. Anim. Behav. Sci.* 59, 331–348.
- Palagi, E., Cordoni, G., Borgognini Tarli, S.M., 2004. Immediate and delayed benefits of play behaviour: New evidence from chimpanzees (*Pan troglodytes*). *Ethology* 110, 949–962.
- Rooney, N.J., Bradshaw, J.W.S., 2002. An experimental study of the effects of play upon the dog-human relationship. *Appl. Anim. Behav. Sci.* 75, 161–176.
- Rooney, N.J., Bradshaw, J.W.S., Robinson, I.H., 2000. A comparison of dog–dog and dog–human play behaviour. *Appl. Anim. Behav. Sci.* 66, 235–248.
- Rooney, N.J., Bradshaw, J.W.S., Robinson, I.H., 2001. Do dogs respond to play signals given by humans? *Anim. Behav.* 61, 715–722.
- Russell, E.S., 1936. Playing with a dog. *Q. Rev. Biol.* 11 (1), 1–15.
- Salimetrics, L.L.C., 2005. *Expanded Range High Sensitivity Salivary Cortisol Enzyme Immunoassay Kit [Brochure]*. State College, PA.
- Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr. Rev.* 21 (1), 55–89.
- Selye, H., 1950. *Stress*. Acta, Inc. Medical Publishers, Montreal, Canada.
- Soderquist, T.R., Serena, M., 2000. Juvenile behaviour and dispersal of chuditch (*Dasyurus geoffroii*) (Marsupialia: Dasyuridae). *Aust. J. Zool.* 48, 551–560.
- Studzinski, C.M., Christie, L.-A., Araujo, J.A., Burnham, W.M., Head, E., Cotman, C.W., Milgram, N.W., 2006. Visuospatial function in the beagle dog: an early marker of cognitive decline in a model of human aging and dementia. *Neurobiol. Learn. Mem.* 86 (2), 197–204.
- Tuber, D.S., Hennessy, M.B., Sanders, S., Miller, J.A., 1996. Behavioral and glucocorticoid responses of adult domestic dogs (*Canis familiaris*) to companionship and social separation. *J. Comp. Psychol.* 110 (1), 103–108.
- Vincent, I.C., Michell, A.R., 1992. Comparison of cortisol concentrations in saliva and plasma of dogs. *Res. Vet. Sci.* 53 (3), 342–345.